

Thank you for taking a few minutes to be involved in our Research Study. Please complete the questionnaire below.

Clicking next will confirm that you have read the Study Participant Information supplied.

* 1. Please enter your date of birth in the format DD/MM/YYYY:

2. What year did you complete your undergraduate medical degree?

3. In what country did you complete your undergraduate medical degree?

4. What type of practice do you currently work in?

- Private General Practice
- Community Health Centre
- Hospital

Other (please specify)

5. Have you ever inserted IUDs before in any setting, either in Australia or overseas?

- Yes No

Give details:

6. What is the main reason you have chosen to do IUD Training at this time?

- Patient demand
- Difficulty in referring on to IUD provider in your area
- Desire for new knowledge and skill
- Financial benefits
- QI and CPD points

Other (please specify)

7. Do any other doctors in your practice currently insert IUDs?

Yes No

Give details:

8. After completing your training, can you see any potential barriers to undertaking IUD insertion in your own practice?

- Time constraints
- Lack of equipment
- Lack of suitable patients
- Level of confidence performing the procedure unsupervised
- Need for Nurse assistance
- Not cost effective

Other (please specify)

9. The statements below reflect knowledge and attitudes towards IUDs. Using the scale please indicate your response to the following statements below:

	Strongly agree	Agree	Undecided	Disagree	Strongly Disagree
Past treatment for an STI is a contraindication to IUD insertion.	<input type="radio"/>				
Screening for Chlamydia is recommended for all women prior to IUD insertion.	<input type="radio"/>				
Screening for Bacterial Vaginosis is recommended for all women prior to IUD insertion.	<input type="radio"/>				
Copper bearing IUDs are a suitable choice for multiparous women.	<input type="radio"/>				
Hormone bearing IUDs (Mirena) are a suitable choice for multiparous women.	<input type="radio"/>				
Copper bearing IUDs are a suitable choice for nulliparous women.	<input type="radio"/>				
Hormone bearing IUDs (Mirena) are a suitable choice for nulliparous women.	<input type="radio"/>				
Copper bearing IUDs can be used for emergency contraception for up to 5 days post unprotected sexual intercourse.	<input type="radio"/>				
Hormone bearing IUDs (Mirena) can be used for emergency contraception for up to 5 days post unprotected sexual intercourse.	<input type="radio"/>				

10. The statements below reflect skills around IUD insertion. Using the scale please indicate your response to the following statements below:

	Strongly agree	Agree	Undecided	Disagree	Strongly Disagree
I have the confidence and skills to insert Copper bearing IUDs in multiparous women.	<input type="radio"/>				
I have the confidence and skills to insert Hormone releasing IUD (Mirena) in multiparous women.	<input type="radio"/>				
I have the confidence and skills to insert Copper bearing IUDs in nulliparous women.	<input type="radio"/>				
I have the confidence and skills to insert Hormone releasing IUD (Mirena) in nulliparous women.	<input type="radio"/>				