



J Simon Bell

Carmel Mezrani

Natalie Blacker

Tammy LeBlanc

Oliver Frank

Christopher P Alderman

Simone Rossi

Debra Rowett

Russell Shute

# Anticholinergic and sedative medicines

## Prescribing considerations for people with dementia

### Background

Older people with dementia may be particularly susceptible to cognitive impairment associated with anticholinergic and sedative medicines. This impairment may be misattributed to the disease process itself.

### Objective

This review examines clinical considerations associated with using anticholinergic and sedative medicines in people with dementia or incipient cognitive impairment. It highlights issues associated with concomitant use of cholinesterase inhibitors and anticholinergic medicines, and pharmacotherapy of conditions that commonly occur in people with dementia.

### Discussion

Use of medicines with anticholinergic or sedative properties may result in adverse events by increasing the overall anticholinergic or sedative load. Patients may benefit from clinicians reviewing the anticholinergic load of the current medicine regimen before the initiation of cholinesterase inhibitors or memantine. Reducing the number and dose of anticholinergic and sedative medicines may improve cognitive function and reduce the likelihood of adverse events.

### Keywords

hypnotics and sedatives; anticholinergic effects; cholinergic antagonists; dementia; cognition disorders

### Medicines with anticholinergic properties

The likelihood that medicines may produce unwanted central anticholinergic effects depends in part on age related and patient specific variability in pharmacokinetic parameters, blood-brain barrier permeability, degree of cholinergic neuronal degeneration and a patient's baseline cognitive status.<sup>4,5</sup> Medicines specifically prescribed for their anticholinergic properties (eg. oxybutynin, benzhexol and benzotropine) are well recognised by clinicians. However, clinicians may be less aware that some medicines prescribed for other purposes also have anticholinergic properties (*Table 1*).<sup>2,6-8</sup> These include the inhaled anticholinergics, ipratropium and tiotropium bromide when prescribed in their usual therapeutic doses. Even medicines with minor anticholinergic properties may contribute to unwanted central and peripheral adverse events if used in combination with other agents with anticholinergic effects. Clinically significant adverse events (*Table 2*) range from mild cognitive impairment to delirium. The cumulative effect of taking one or more medicines with anticholinergic properties has been termed 'anticholinergic load'.<sup>9</sup>

Use of medicines with anticholinergic properties in older people is associated with impaired physical and cognitive function. In a cross sectional study of 364 people aged more than 80 years living in Italy, the use of medicines with serum anticholinergic activity was associated with poorer physical performance battery scores, reduced hand grip strength and compromised activities of daily living.<sup>10</sup> In 932 community dwelling people aged more than 65

The number of people with dementia in Australia is predicted to quadruple from an estimated 245 000 in 2009 to approximately 1.13 million by 2050.<sup>1</sup> Causes of dementia include Alzheimer disease, dementia with Lewy bodies, frontotemporal dementia, vascular dementia and Parkinson disease. Medicines with anticholinergic and sedative properties are widely prescribed for older people in Australia.<sup>2,3</sup> People with dementia may be particularly susceptible to cognitive impairment caused by anticholinergic and sedative medicines.<sup>4</sup>

**Table 1. Medicines with clinically significant anticholinergic effects that are commonly used in older people with dementia<sup>6–8</sup>**

	Antipsychotics	Antidepressants	Medicines for urinary incontinence	Antihistamines
Strong anticholinergic effects – avoid using in people with dementia	Chlorpromazine Olanzapine Pericyazine	Tricyclic antidepressants (eg. amitriptyline, doxepin, imipramine)	Darifenacin** Oxybutynin Propantheline Solifenacin** Tolterodine**	Brompheniramine*** Chlorpheniramine*** Cyproheptadine Diphenhydramine*** Promethazine***
Moderate anticholinergic effects – use with caution in people with dementia	Haloperidol Prochlorperazine Quetiapine Risperidone Ziprasidone	Desvenlafaxine Duloxetine* Fluoxetine Mirtazapine Paroxetine Reboxetine* Venlafaxine		

\* New medicine: reported adverse effects profile is consistent with moderate anticholinergic effects

\*\* Not included on the schedule of PBS/RPBS benefits

\*\*\* Found in cold and flu treatments, may be purchased over-the-counter

Reproduced from Veterans' MATES Therapeutic brief 25. Available at [www.veteransmates.net.au](http://www.veteransmates.net.au)

**Table 2. Examples of anticholinergic side effects**

Confusion/hallucinations/delirium  
Dry mouth  
Pupil dilatation/blurred vision  
Urinary retention  
Constipation  
Tachycardia/arrhythmias

Reproduced from Veterans' MATES Therapeutic brief 25. Available at [www.veteransmates.net.au](http://www.veteransmates.net.au)

also have sedative properties. Medicines with sedative properties have been linked to depressive symptoms, worsening cognition, respiratory depression, impaired muscle strength and falls and fractures.<sup>15–19</sup> The cumulative effect of taking one or more medicines with sedative properties has been termed 'sedative load'.<sup>20</sup> All medicines with sedative properties – not only those prescribed for intentional sedation – may contribute to an older person's sedative load.

### Concomitant use of cholinesterase inhibitors and anticholinergic medicines

Cholinesterase inhibitors (donepezil, rivastigmine and galantamine) may modestly improve cognition via inhibiting the enzyme acetyl cholinesterase. Restricted supply of these medicines is subsidised through the Pharmaceutical Benefits Scheme (PBS) using the authority system for the management of mild to moderate Alzheimer disease. Concomitant use of anticholinergic medicines may decrease the effectiveness of cholinesterase inhibitors.<sup>21</sup> Despite this, an Australian study revealed that of 5797 people who commenced treatment with a PBS subsidised cholinesterase inhibitor between April and June 2006, 32% also received a prescription for an anticholinergic medicine in the 14 weeks before

and following the first date of supply.<sup>3</sup> Dispensing of anticholinergic medicines also increased following initiation of cholinesterase inhibitors. Some of these anticholinergic medicines may have been prescribed to treat the adverse effects of the cholinesterase inhibitor (eg. medicines for urinary incontinence). Patients may benefit from clinicians reviewing the anticholinergic load of their current medicine regimen before initiating cholinesterase inhibitors or memantine, with the aim of minimising or ceasing medicines with anticholinergic properties.

### Reducing anticholinergic and sedative load in patients with dementia

#### Pharmacotherapy of specific conditions

##### Urinary incontinence

Incontinence is often multifactorial in older patients with dementia. Medicines may contribute to urinary incontinence via agonism of alpha-1-adrenoceptors or nicotinic acid receptors leading to stress incontinence, or antagonism of beta-3-adrenoceptors or agonism of muscarinic receptors leading to urge incontinence.<sup>22</sup> Antagonism of muscarinic receptors may lead to overflow incontinence, while use of medicines with

years in the Women's Health and Aging Study conducted in the United States, use of medicines with anticholinergic properties was associated with difficulties in balance and mobility.<sup>11</sup> There is mixed evidence in relation to an association between anticholinergic medicines and increased mortality.<sup>12–14</sup>

### Medicines with sedative properties

Medicines with sedative properties include benzodiazepines and other hypnotics, antipsychotics, anticonvulsants, antidepressants, opioid analgesics and tramadol, and histamine H1 receptor antagonists commonly used for allergic conditions. Many anticholinergic medicines

sedative properties (eg. benzodiazepines, opioids and tricyclic antidepressants) may contribute to functional incontinence. Anticholinergic medicines can cause constipation, which can result in urinary retention and urge and overflow incontinence.

Cholinesterase inhibitors prescribed for patients with Alzheimer disease have also been associated with urinary incontinence.<sup>23</sup> In a retrospective population based cohort study of 44 884 older people with dementia, use of cholinesterase inhibitors was associated with an increased likelihood of receiving an anticholinergic medicine to manage urinary incontinence.<sup>24</sup> Drug induced urinary incontinence in people with dementia may be misattributed to progression of the underlying disease processes.<sup>25</sup> This may result in a ‘prescribing cascade’ if an anticholinergic medicine is then prescribed to treat the incontinence.<sup>24</sup>

Nonpharmacological approaches are recommended as first line treatment. Prompted or timed voiding may reduce urinary incontinence among residents of aged care facilities with dementia. For people with functional incontinence related to impaired mobility, an occupational therapist or physiotherapist may advise measures to improve toilet access (eg. removal of clutter, good lighting, nonslip flooring, ensuring the toilet is clearly marked, hand rails, raised toilet seat height and appropriate clothing). With these interventions, treatment with anticholinergic medicines may not be necessary. However, if anticholinergic medicines are to be trialled, patients and carers may require education about potential adverse events of anticholinergic medicines. Consider ceasing anticholinergics administered for urinary symptoms if there is no benefit after 4 weeks of treatment.<sup>8</sup>

## Depression

Depressive symptoms have been reported in up to 40% of people with Alzheimer disease, which is reflected in the high prevalence of antidepressant use in this population.<sup>26</sup> However, there is a lack of evidence to guide antidepressant prescribing.<sup>27</sup> Two recent studies have suggested a less favourable risk-to-benefit ratio than previously thought. An observational study reported an increased risk of falls and fractures, stroke/transient ischaemic attack and all cause mortality among users of selective serotonin

reuptake inhibitors (SSRIs) compared to tricyclic antidepressants (TCAs).<sup>28</sup> The Health Technology Assessment Study of the Use of Antidepressants for Depression in Dementia (HTA-SADD), which included 326 patients, reported that sertraline and mirtazapine were not more effective than placebo and were associated with an increased risk of adverse events.<sup>29</sup> This finding contradicted an earlier meta-analysis that included 165 patients which supported the efficacy of antidepressants for treating depression in Alzheimer disease.<sup>30</sup> A Cochrane review has also reported that sertraline and citalopram were associated with a reduction in agitation in people with dementia when compared to placebo in two studies.<sup>31</sup> Selective serotonin reuptake inhibitors tend to be less sedating than TCAs, mianserin or mirtazapine. Some TCAs (eg. amitriptyline, doxepin) are both highly sedative and anticholinergic. Combination antidepressant treatment is not appropriate due to the increased risk of adverse events, including those related to anticholinergic and sedative load.<sup>32</sup>

## Insomnia

Sleep complaints are common among people with Alzheimer disease and other dementias.<sup>33</sup> Underlying issues that may contribute to disturbed sleep include medicines, medical conditions and environmental factors. Older people often require less sleep than younger people and it is useful to explain this to patients. Nonpharmacological treatments for insomnia are considered first line.<sup>34</sup> Sedative-hypnotic medicines should be prescribed for the shortest possible duration as an adjunct to nonpharmacological treatments. It is useful to agree a definite duration of treatment with the patient at the time of prescribing.<sup>8</sup> People already taking long term benzodiazepines may benefit from dosage reduction and slow discontinuation to maximise cognitive function and reduce the risk of falls. Despite their relatively widespread use, TCAs and sedating antihistamines are best avoided in the management of insomnia due to their anticholinergic properties.

## Pain management

Opioid analgesics and tramadol may have sedative effects. The ways in which people with dementia experience and express pain is a field of ongoing research.<sup>35</sup> People with dementia may

express pain in the form of facial expressions, body movements and behavioural disturbances such as aggression or agitation. These symptoms may be misinterpreted as behavioural symptoms of dementia. Pain may be under-recognised and undertreated, especially in advanced stages of dementia.<sup>36</sup> Research has reported lower overall use of analgesics among people with dementia.<sup>37,38</sup> However, while people with dementia may receive less analgesics overall, they may be more likely to be prescribed strong opioids (eg. transdermal fentanyl) instead of paracetamol and other analgesics with lower likelihoods of adverse events.<sup>38</sup> Transdermal fentanyl patches should not be prescribed to people with chronic nonmalignant pain who are opioid naive.<sup>8</sup> Antiepileptics used in the treatment of neuropathic pain (eg. gabapentin, pregabalin, carbamazepine, valproate) may impair cognition and cause sedation. People with dementia may also be particularly susceptible to memory impairment and cognitive decline associated with some antiepileptics.<sup>39</sup>

## Behavioural and psychological symptoms

Behavioural and psychological symptoms of dementia are common and include agitation, aggression, hallucinations and wandering. Nonpharmacological strategies are considered first line treatment. Treatment with antipsychotics should be reserved for people with aggression and psychosis who have not responded to nonpharmacological strategies. Use of both first (eg. haloperidol) and second generation (eg. olanzapine, quetiapine, risperidone) antipsychotics is associated with an increased risk of death in people with Alzheimer disease.<sup>40,41</sup> In common with some other medicines commonly prescribed for older people (eg. SSRIs, TCAs, opioids) antipsychotics can also cause hyponatraemia that may result in confusion and lethargy. While second generation antipsychotics have a lower risk of tardive dyskinesia, they can cause sedation and postural hypotension. Patients with Lewy body dementia are especially susceptible to the extrapyramidal side effects of antipsychotics.<sup>42</sup> The dementia antipsychotic withdrawal trial (DART-AD) in the United Kingdom demonstrated that for most patients with Alzheimer disease,

withdrawal of antipsychotics had no detrimental effect on functional and cognitive status. People who continued using antipsychotics had reduced survival compared to those who received placebo (ie. those who discontinued antipsychotics treatment).<sup>43</sup> In Australia, risperidone is the only antipsychotic PBS listed for the management of behavioural disturbances characterised by psychotic symptoms and aggression in patients with dementia where nonpharmacological methods have been unsuccessful. Problems such as screaming/vocalising, ‘sundowning’ and wandering do not reliably respond to antipsychotics.

## Key points

- Older people with dementia may be particularly susceptible to impaired physical and cognitive function associated with anticholinergic and sedative medicines.
- Drug induced physical and cognitive impairment in older people with dementia may be misattributed to progression of underlying disease processes.
- Medicines with weak anticholinergic or sedative properties (including nonprescription medicines) may cause adverse events by contributing to an older person’s anticholinergic or sedative load. This load may be decreased by reducing the number and dose of medicines with anticholinergic and sedative properties.
- Patients may benefit from clinicians reviewing the anticholinergic load of their current medicine regimen before initiating cholinesterase inhibitors or memantine, with the aim of minimising or ceasing if possible medicines with anticholinergic properties.
- Urinary incontinence may be drug induced. Anticholinergic medicines are often not needed to treat urinary incontinence. Consider ceasing anticholinergic medicines if there is no benefit after 4 weeks.
- Depressive symptoms are common in Alzheimer disease. There is mixed evidence for the safety and efficacy of antidepressants in patients with dementia. Selective serotonin reuptake inhibitors are less sedating than TCAs, mianserin or mirtazapine. Tricyclic antidepressants may be both sedative and anticholinergic.

- For most patients with Alzheimer disease, withdrawal of antipsychotics has no detrimental effect on functional and cognitive status.

## Authors

J Simon Bell BPharm(Hons), PhD, MPS, is Associate Professor, Quality Use of Medicines and Pharmacy Research Centre, Sansom Institute, School of Pharmacy and Medical Sciences, University of South Australia. [simon.bell@unisa.edu.au](mailto:simon.bell@unisa.edu.au)

Carmel Mezrani BMedSc, BMBS, FRACGP, is Senior Lecturer, School of Medicine, University of Notre Dame, Sydney, New South Wales and the Department of Veterans’ Affairs, Veterans’ Medicines Advice and Therapeutics Education Services (Veterans’ MATES) Clinical Reference Group

Natalie Blacker BBehavSc(Psych), is Research Associate, Quality Use of Medicines and Pharmacy Research Centre, Sansom Institute, School of Pharmacy and Medical Sciences, University of South Australia

Tammy LeBlanc BA(Psych), is module development coordinator, Quality Use of Medicines and Pharmacy Research Centre, Sansom Institute, School of Pharmacy and Medical Sciences, University of South Australia

Oliver Frank MBBS, PhD, FRACGP, FACHI, is University Senior Research Fellow, Discipline of General Practice, School of Population Health and Clinical Practice, The University of Adelaide, South Australia and the Department of Veterans’ Affairs, Veterans’ Medicines Advice and Therapeutics Education Services (Veterans’ MATES) Clinical Reference Group

Christopher P Alderman BPharm, PhD, FSHP, CGP, BCPP, is Associate Professor, Pharmacy Department, Repatriation General Hospital, Adelaide, South Australia and the Department of Veterans’ Affairs, Veterans’ Medicines Advice and Therapeutics Education Services (Veterans’ MATES) Clinical Reference Group

Simone Rossi BPharm, is Editor, Australian Medicines Handbook, Adelaide, South Australia and the Department of Veterans’ Affairs, Veterans’ Medicines Advice and Therapeutics Education Services (Veterans’ MATES) Clinical Reference Group

Debra Rowett BPharm, is Service Director, Drug and Therapeutics Information Service, Repatriation General Hospital, Adelaide, South Australia and the Department of Veterans’ Affairs, Veterans’ Medicines Advice and Therapeutics Education Services (Veterans’ MATES) Clinical Reference Group

Russell Shute BMBS, ClinDipPaIIMed, FRACGP, is a general practitioner, Australian Medicines Handbook, Adelaide, South Australia and the Department of Veterans’ Affairs, Veterans’ Medicines Advice and Therapeutics Education Services (Veterans’ MATES) Clinical Reference Group.

Conflict of interest: All authors have completed the ICMJE disclosure form of potential conflict of interest. Oliver Frank is employed by the University of Adelaide which has a contract to provide GP advice to the DVA MATES program. Christopher Alderman has received payment from various pharmaceutical companies for lectures, consultancy, expert testimony and for the development of educational presentations. Simone Rossi is employed by the National Prescribing Service Ltd. Debra Rowett is employed by the Repatriation General Hospital which receives funding from DVA.

## Disclaimer

This work has been produced with the assistance of funding provided by the Australian Government Department of Veterans’ Affairs. However, the views expressed do not necessarily represent the views of the Minister for Veterans’ Affairs or the Department of Veterans’ Affairs. The Commonwealth does not give any warranty nor accept any liability in relation to the contents of this work.

## Acknowledgements

The Department of Veterans’ Affairs’ MATES program is provided by the Quality Use of Medicines and Pharmacy Research Centre, Sansom Institute, University of South Australia in association with Discipline of General Practice, University of Adelaide; Discipline of Public Health, University of Adelaide; Repatriation General Hospital, Daw Park; National Prescribing Service; Australian Medicines Handbook; and Drug and Therapeutics Information Service. Veterans’ MATES Program materials are available at [www.veteransmates.net.au](http://www.veteransmates.net.au).

## References

1. Access Economics. Keeping dementia front of mind: incidence and prevalence 2009–2050. 2009. Available at [www.alzheimers.org.au/common/files/NAT/20090800\\_Nat\\_AE\\_FullKeepDemFrontMind.pdf](http://www.alzheimers.org.au/common/files/NAT/20090800_Nat_AE_FullKeepDemFrontMind.pdf) [Accessed 31 May 2011].
2. Nishtala PS, Fois RA, McLachlan AJ, et al. Anticholinergic activity of commonly prescribed medications and neuropsychiatric adverse events in older people. *J Clin Pharmacol* 2009;49:1176–84.
3. Robinson M, Rowett D, Leverton A, et al. Changes in utilisation of anticholinergic drugs after initiation of cholinesterase inhibitors. *Pharmacoepidemiol Drug Saf* 2009;18:659–64.
4. Sunderland T, Tariot PN, Cohen RM, et al.

- Anticholinergic sensitivity in patients with dementia of the Alzheimer type and age-matched controls. A dose-response study. *Arch Gen Psychiatry* 1987;44:418–26.
5. Roe CM, Anderson MJ, Spivack B. Use of anticholinergic medications by older adults with dementia. *J Am Geriatr Soc* 2002;50:836–42.
  6. Rudolph JL, Salow MJ, Angelini MC, et al. The anticholinergic risk scale and anticholinergic adverse effects in older people. *Arch Intern Med* 2008;168:508–13.
  7. Carnahan RM, Lund BC, Perry PJ, et al. The Anticholinergic Drug Scale as a measure of drug-related anticholinergic burden: associations with serum anticholinergic activity. *J Clin Pharmacol* 2006;46:1481–6.
  8. Rossi S, editor. *Australian Medicines Handbook*. Adelaide: Australian Medicines Handbook Pty Ltd; 2011.
  9. Minzenberg MJ, Poole JH, Benton C, et al. Association of anticholinergic load with impairment of complex attention and memory in schizophrenia. *Am J Psychiatry* 2004;161:116–24.
  10. Landi F, Russo A, Liperoti R, et al. Anticholinergic drugs and physical function among frail elderly population. *Clin Pharmacol Ther* 2007;81:235–41.
  11. Cao YJ, Mager DE, Simonsick EM, et al. Physical and cognitive performance and burden of anticholinergics, sedatives, and ACE inhibitors in older women. *Clin Pharmacol Ther* 2008;83:422–9.
  12. Kumpula EK, Bell JS, Soini H, et al. Anticholinergic drug use and mortality among residents of long-term care facilities: a prospective cohort study. *J Clin Pharmacol* 2011;51:256–63.
  13. Uusvaara J, Pitkala KH, Kautiainen H, et al. Association of anticholinergic drugs with hospitalization and mortality among older cardiovascular patients: A prospective study. *Drugs Aging* 2011;28:131–8.
  14. Panula J, Puustinen J, Jaatinen P, et al. Effects of potent anticholinergics, sedatives and antipsychotics on postoperative mortality in elderly patients with hip fracture: a retrospective, population-based study. *Drugs Aging* 2009;26:963–71.
  15. van Vliet P, van der Mast RC, van den Broek M, et al. Use of benzodiazepines, depressive symptoms and cognitive function in old age. *Int J Geriatr Psychiatry* 2009;24:500–8.
  16. Hartikainen S, Lönnroos E, Louhivuori K. Medication as a risk factor for falls: critical systematic review. *J Gerontol A Biol Sci Med Sci* 2007;62:1172–81.
  17. Pariante A, Dartigues JF, Benichou J, et al. Benzodiazepines and injurious falls in community dwelling elders. *Drugs Aging* 2008;25:61–70.
  18. Roth T. Hypnotic use for insomnia management in chronic obstructive pulmonary disease. *Sleep Med* 2009;10:19–25.
  19. Taipale HT, Bell JS, Gnjdic D, et al. Muscle strength and sedative load in community-dwelling people aged 75 years and older: a population-based study. *J Gerontol A Biol Sci Med Sci* 2011; Epub Sept 20 DOI 10.1093/gerona/glr170.
  20. Taipale HT, Bell JS, Soini H, et al. Sedative load and mortality among residents of long-term care facilities: a prospective cohort study. *Drugs Aging* 2009;26:871–81.
  21. Johnell K, Fastbom J. Concurrent use of anticholinergic drugs and cholinesterase inhibitors: register-based study of over 700,000 elderly patients. *Drugs Aging* 2008;25:871–7.
  22. Tsakiris P, Oelke M, Michel MC. Drug-induced urinary incontinence. *Drugs Aging* 2008;25:541–9.
  23. Starr JM. Cholinesterase inhibitor treatment and urinary incontinence in Alzheimer's disease. *J Am Geriatr Soc* 2007;55:500–1.
  24. Gill SS, Mamdani M, Naglie G, et al. A prescribing cascade involving cholinesterase inhibitors and anticholinergic drugs. *Arch Intern Med* 2005;165:808–13.
  25. Resnick NM. Urinary incontinence. *Lancet* 1995;346:94–9.
  26. Arbus C, Gardette V, Bui E, et al. Antidepressant use in Alzheimer's disease patients: results of the REAL FR cohort. *Int Psychogeriatr* 2010;22:120–8.
  27. Nelson JC, Devanand DP. A systematic review and meta-analysis of placebo-controlled antidepressant studies in people with depression and dementia. *J Am Geriatr Soc* 2011;59:577–85.
  28. Coupland C, Dhiman P, Morriss R, et al. Antidepressant use and risk of adverse outcomes in older people: population based cohort study. *BMJ* 2011;343:d4551.
  29. Banerjee S, Hellier J, Dewey M, et al. Sertraline or mirtazapine for depression in dementia (HTA-SADD): a randomised, multicentre, double-blind, placebo-controlled trial. *Lancet* 2011;378:403–11.
  30. Thompson S, Herrmann N, Rapoport MJ, et al. Efficacy and safety of antidepressants for treatment of depression in Alzheimer's disease: a metaanalysis. *Can J Psychiatry* 2007;52:248–55.
  31. Seitz DP, Adunuri N, Gill SS, et al. Antidepressants for agitation and psychosis in dementia. *Cochrane Database Syst Rev* 2011;2:CD008191.
  32. Rush AJ, Trivedi MH, Stewart JW, et al. Combining medications to enhance depression outcomes (CO-MED): acute and long-term outcomes of a single-blind randomized study. *Am J Psychiatry* 2011;168:689–701.
  33. Bombois S, Derambure P, Pasquier F, et al. Sleep disorders in aging and dementia. *J Nutr Health Aging* 2010;14:212–7.
  34. David R, Zeitzer J, Friedman L, et al. Non-pharmacologic management of sleep disturbance in Alzheimer's disease. *J Nutr Health Aging* 2010;14:203–6.
  35. Scherder E, Herr K, Pickering G, et al. Pain in dementia. *Pain* 2009;145:276–8.
  36. Morrison RS, Siu AL. A comparison of pain and its treatment in advanced dementia and cognitively intact patients with hip fracture. *J Pain Symptom Manage* 2000;19:240–8.
  37. Nygaard HA, Jarland M. Are nursing home patients with dementia diagnosis at increased risk for inadequate pain treatment? *Int J Geriatr Psychiatry* 2005;20:730–7.
  38. Bell JS, Laitinen ML, Lavikainen P, et al. Use of strong opioids among community-dwelling persons with and without Alzheimer's disease in Finland. *Pain* 2011;152:543–7.
  39. Kwan P, Brodie MJ. Neuropsychological effects of epilepsy and antiepileptic drugs. *Lancet* 2001;357:216–22.
  40. Setoguchi S, Wang PS, Brookhart MA, et al. Potential causes of higher mortality in elderly users of conventional and atypical antipsychotic medications. *J Am Geriatr Soc* 2008;56:1644–50.
  41. Wang PS, Schneeweiss S, Avorn J, et al. Risk of death in elderly users of conventional vs. atypical antipsychotic medications. *N Engl J Med* 2005;353:2335–41.
  42. Weintraub D, Hurtig HI. Presentation and management of psychosis in Parkinson's disease and dementia with Lewy bodies. *Am J Psychiatry* 2007;164:1491–8.
  43. Ballard C, Hanney ML, Theodoulou M, et al. The dementia antipsychotic withdrawal trial (DART-AD): long-term follow-up of a randomised placebo-controlled trial. *Lancet Neurol* 2009;8:151–7.

correspondence [afp@racgp.org.au](mailto:afp@racgp.org.au)